

## Agrobacterium articles

Coutu, C., Brandle, J., Brown, D., Brown, K., Miki, B., Simmonds, J., & Hegedus, D. (2007). PORE: A modular binary vector series suited for both monocot and dicot plant transformation. *Transgenic Res Transgenic Research*, 771-781. doi:10.1007/s11248-007-9066-2

Gelvin, S. (2012). Traversing the Cell: Agrobacterium T-DNA's Journey to the Host Genome. *Frontiers in Plant Science Front. Plant Sci.* doi:10.3389/fpls.2012.00052

Bell, P. (2001). Ecoli Expression Systems (A. Gerstein, Ed.). *Molecular Biology Problem Solver: A Laboratory Guide*, 461-490. doi:2001

Hellens, R., Mullineaux, P., & Klee, H. (2000). Technical Focus: A guide to Agrobacterium binary Ti vectors. *Trends in Plant Science*, 5(10), 446-451.

Kiyokawa, K., Yamamoto, S., Sakuma, K., Tanaka, K., Moriguchi, K., & Suzuki, K. (2009). Construction of Disarmed Ti Plasmids Transferable between Escherichia coli and Agrobacterium Species. *APPLIED AND ENVIRONMENTAL MICROBIOLOGY*, 1845-1851.

Lacroix, B., Tzfira, T., Vainstein, A., & Citovsky, V. (n.d.). A case of promiscuity: Agrobacterium's endless hunt for new partners. *Trends in Genetics*, 29-37.

Riva, G., Gonzalez-Cabrera, J., Vazquez-Padron, R., & Ayra-Pardo, C. (1998). Agrobacterium tumefaciens : A natural tool for plant transformation. *Electron. J. Biotechnol. Electronic Journal of Biotechnology*, 118-113.

Tinland, B., Hohn, B., & Puchta, H. (1994). Agrobacterium tumefaciens transfers single-stranded transferred DNA (T-DNA) into the plant cell nucleus. *Proceedings of the National Academy of Sciences*, 91, 8000-8004.

Wise, A., Liu, Z., & Binns, A. (2006). Culture and Maintenance of Agrobacterium Strains. *Agrobacterium Protocols*, 3-14.

Yadav, N., Vanderleyden, J., Bennett, D., Barnes, W., & Chilton, M. (1982). Short direct repeats flank the T-DNA on a nopaline Ti plasmid. *Proceedings of the National Academy of Sciences*, 79, 6322-6326.

## Cannabidiol Article

Shoyama, Y., Tamada, T., Kurihara, K., Takeuchi, A., Taura, F., Arai, S., . . . Kuroki, R. (n.d.). Structure and Function of  $\Delta^1$ -Tetrahydrocannabinolic Acid (THCA) Synthase, the Enzyme Controlling the Psychoactivity of Cannabis sativa. *Journal of Molecular Biology*, 96-105.

Sirikantaramas, B., Taura, B., Morimoto, B., & Shoyama, B. (2007). Recent Advances in Cannabis sativa Research: Biosynthetic Studies and Its Potential in Biotechnology. *Current Pharmaceutical Biotechnology CPB*, 237-243.

Sirikantaramas, S., Morimoto, S., Shoyama, Y., Ishikawa, Y., Wada, Y., Shoyama, Y., & Taura, F. (2004). The Gene Controlling Marijuana Psychoactivity: MOLECULAR CLONING AND HETEROLOGOUS EXPRESSION OF 1-TETRAHYDROCANNABINOLIC ACID SYNTHASE FROM CANNABIS SATIVA L. *Journal of Biological Chemistry*, 39767-39774.

Sirikantaramas, S. (2005). Tetrahydrocannabinolic Acid Synthase, the Enzyme Controlling Marijuana Psychoactivity, is Secreted into the Storage Cavity of the Glandular Trichomes. *Plant and Cell Physiology*, 1578-1582. doi:10.1093/pcp/pci166,

Takeda, S., Okajima, S., Miyoshi, H., Yoshida, K., Okamoto, Y., Okada, T., ... & Aramaki, H. (2012). Cannabidiolic acid, a major cannabinoid in fiber-type cannabis, is an inhibitor of MDA-MB-231 breast cancer cell migration. *Toxicology letters*, 214(3), 314-319.

Taura, F., Sirikantaramas, S., Shoyama, Y., Yoshikai, K., Shoyama, Y., & Morimoto, S. (n.d.). Cannabidiolic acid synthase, the chemotype-determining enzyme in the fiber-type Cannabis sativa. *FEBS Letters*, 2929-2934.

## Mambalgin Articles

Matsumoto, M. (2013). Faculty of 1000 evaluation for Black mamba venom peptides target acid-sensing ion channels to abolish pain. *F1000 - Post-publication Peer Review of the Biomedical Literature*. doi:10.1038/nature11494

Sherwood, T., Frey, E., & Askwith, C. (2012). Structure and activity of the acid-sensing ion channels. *AJP: Cell Physiology*, C699-C710. doi:10.1152/ajpcell.00188.2012

Zha, X. (2013). Acid-sensing ion channels: Trafficking and synaptic function. *Molecular Brain Mol Brain*, 1-1.

## Pichia Articles

Ahmad, M., Hirz, M., Pichler, H., & Schwab, H. (2014). Protein expression in Pichia pastoris: Recent achievements and perspectives for heterologous protein production. *Appl Microbiol Biotechnol Applied Microbiology and Biotechnology*, 5301-5317. doi:10.1007/s00253-014-5732-5

Altstein, M. (2004). Peptide pheromones: An overview. *Peptides*, 1373-1376.

Basanta, A., Gomez-Sala, B., Sanchez, J., Diep, D., Herranz, C., Hernandez, P., & Cintas, L. (2010). Use of the Yeast Pichia pastoris as an Expression Host for Secretion of Enterocin L50, a Leaderless Two-Peptide (L50A and L50B) Bacteriocin from Enterococcus faecium L50. *APPLIED AND ENVIRONMENTAL MICROBIOLOGY*, 76(10), 3314-3324. doi:10.1128/AEM.02206-09

Cereghino, J. (1999). Heterologous protein expression in the methylotrophic yeast *Pichia pastoris*. *FEMS Microbiology Reviews*, 45-66.

James, C. (2007). *Pichia Protocols*, Second Edition.

Kawai, S., Hashimoto, W., & Murata, K. (n.d.). Transformation of *Saccharomyces cerevisiae* and other fungi. *Bioengineered Bugs*, 395-403.

Plessis, A., & Dujon, B. (1993). Multiple tandem integrations of transforming DNA sequences in yeast chromosomes suggest a mechanism for integrative transformation by homologous recombination. *Gene*, 41-50.

Qin, X., Qian, J., Yao, G., Zhuang, Y., Zhang, S., & Chu, J. (2011). GAP Promoter Library for Fine-Tuning of Gene Expression in *Pichia pastoris*. *APPLIED AND ENVIRONMENTAL MICROBIOLOGY*, 77(11), 3600-3608. doi:10.1128/AEM.02843-10